

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/50939628>

Continuous Transcutaneous Submental Electrical Stimulation in Obstructive Sleep Apnea A Feasibility Study

Article in *Chest* · March 2011

DOI: 10.1378/chest.10-2614 · Source: PubMed

CITATIONS

22

READS

137

9 authors, including:



John Seymour

Frimley Park Hospital NHS Foundation Trust

39 PUBLICATIONS 1,590 CITATIONS

[SEE PROFILE](#)



Gerrard Rafferty

King's College Hospital NHS Foundation Tr...

230 PUBLICATIONS 3,479 CITATIONS

[SEE PROFILE](#)



Caroline J Jolley

King's College London

86 PUBLICATIONS 1,457 CITATIONS

[SEE PROFILE](#)



Yuanming Luo

State Key Laboratory of Respiratory Medicine

105 PUBLICATIONS 1,529 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Maternal Medicine [View project](#)



SIDS and CDH [View project](#)



Continuous Transcutaneous Submental Electrical Stimulation in Obstructive Sleep Apnea

A Feasibility Study

Joerg Steier, MD, PhD; John Seymour, PhD; Gerrard F. Rafferty, PhD;
Caroline J. Jolley, MD; Eskinder Solomon, MSc; Yuanming Luo, PhD;
William D.-C. Man, PhD; Michael I. Polkey, PhD; and John Moxham, MD

Background: The therapeutic value of transcutaneous electrical stimulation of the genioglossus muscle in patients with obstructive sleep apnea (OSA) to reduce sleep-disordered breathing is unclear.

Methods: Contraction of the genioglossus muscles during transcutaneous stimulation was investigated using ultrasonography in 11 healthy subjects (seven men, mean [SD] age 30 [6] years; BMI, 24.2 [3.5] kg/m²). Esophageal and gastric pressures were measured with balloon catheters, and transesophageal diaphragm electromyogram (EMGdi) was recorded during polysomnography in 11 patients with OSA (eight men, aged 51 [16] years; BMI, 42.0 [9.7] kg/m²) while transcutaneous electrical stimulation of the genioglossus was applied in non-rapid eye movement sleep (stage N2).

Results: Ultrasonography measurements showed a significant increase in tongue diameter during stimulation (sagittal: 10.0% [2.8%]; coronal: 9.4% [3.7%]). The measurements were reproducible and repeatable. In patients with OSA, snoring decreased during stimulation ($P < .001$) and oxygenation improved ($P = .001$); the respiratory disturbance index (RDI) fell from 28.1 (26.3) to 10.2 (10.2) events per hour during stimulation ($P = .002$), returning to 26.6 (26.0) events per hour after stimulation was stopped. Transdiaphragmatic pressure swing decreased from 24.1 (13.5) cm H₂O to 19.7 (7.1) cm H₂O ($P = .022$), increasing to 24.2 (10.8) cm H₂O afterward, and EMGdi fell from 23.8% max (12.6% max) to 15.7% max (6.4% max) ($P < .001$), rising to 22.6% max (10.4% max) post stimulation.

Conclusions: Continuous transcutaneous electrical stimulation of the genioglossus contracts the genioglossus muscle and reduces ventilatory load and neural respiratory drive in patients with OSA.

CHEST 2011; 140(4):998–1007

Abbreviations: AHI = apnea-hypopnea index; CPAP = continuous positive airway pressure; Ds-t = distance skin-to-tongue surface; EMG = electromyogram; EMGdi = transesophageal diaphragm electromyogram; OSA = obstructive sleep apnea; Pdi = transdiaphragmatic pressure; Pes = esophageal pressure; RDI = respiratory disturbance index; SaO₂ = oxygen saturation

Obstructive sleep apnea (OSA) is a common disorder among middle-aged adults, affecting approximately 4% of men and 2% of women.¹ Continuous positive airway pressure (CPAP) during sleep is the best available treatment² for the majority of patients in addition to appropriate lifestyle interventions.²⁻⁶ However, some patients may not comply with or respond to CPAP; approximately one-third of all patients initiated on CPAP do not use the treatment at 5 years.⁷

Remmers et al⁸ hypothesized that upper airway occlusion in sleep could be countered by activity of

the dilators of the upper airway. This hypothesis has been tested by studying the dilator muscles,⁹⁻¹⁵ in particular the genioglossus muscle,¹⁶⁻¹⁸ by electrically stimulating them using intraoral surface or intramuscular fine wire electrodes. However, it is difficult to adequately stimulate sufficient bulk of the pharyngeal dilator muscles, and such stimulation often disturbs sleep.

Transcutaneous stimulation of the pharyngeal dilators, in particular the genioglossus, may represent an alternative and better-tolerated approach. However,

conflicting data exist as to the efficacy of this method.^{19,20} Miki et al¹⁹ demonstrated that short periods of transcutaneous electrical stimulation decreased the incidence of episodes of upper airway occlusion in OSA. However, Edmonds et al²⁰ found transcutaneous electrical stimulation to be ineffective. We hypothesized that low-current transcutaneous electrical stimulation, applied over a longer period, could result in sufficient contraction of the genioglossus muscle to reduce sleep-disordered breathing in patients with OSA without awakening.

MATERIALS AND METHODS

The first part of the study sought to determine a suitable stimulation frequency and intensity that was tolerable and to evaluate with ultrasonography whether transcutaneous electrical stimulation of the genioglossus achieves an effective muscle contraction. Next, we observed the response to stimulation during sleep and studied whether respiratory effort in OSA was reduced.

We used ultrasonography to investigate genioglossus muscle responses to transcutaneous stimulation in 11 awake normal subjects, then studied the effect of stimulation on sleep-disordered breathing in 11 subjects with known OSA. Participants were initially addressed following referral from the metabolic clinic at King's College Hospital. Patients were excluded if they had significant comorbidities, particularly lung disease. We did not include patients who had been treated for lung disease such as asthma or obesity-hypoventilation syndrome previously. In addition, we did not recruit from patients with acute illness. Eight of our patients with sleep apnea had diabetes, but this was well controlled at the time of the study. The study was approved by King's College Hospital local research ethics committee (reference number 06/Q0703/224), and each participant gave informed written consent.

Transcutaneous Electrical Stimulation of Genioglossus

The skin was prepared with alcohol wipes and Nuprep skin gel (D. O. Weaver Co; Aurora, Colorado). Two patches (40 × 40 mm) (Verity Medical Ltd; Uplands, England) were placed halfway between the chin and the angle of the mandible over the submental area (Figs 1A, 1B) to deliver transcutaneous stimulation with a NeuroTrac Sports stimulator (Verity Medical Ltd). Stimulation intensity, at a frequency of 30 Hz for approximately 1 min, was

titrated to ensure it was comfortable while the patient was awake. Having established a comfortable level of stimulation, the maximal tolerable level of current was also determined.

Ultrasonography of the Genioglossus

Ultrasonography was used to measure the thickness of the tongue muscles with an 8-MHz, 5.6-cm linear probe (Toshiba Medical Systems; Tokyo, Japan). Images were taken in both the sagittal and coronal planes with the probe placed under the mandible (Figs 2A, 2B) (additional information on how these measurements were made is available in e-Appendix 1).

The measurement of the distance skin-to-tongue surface (Ds-t) was taken during electrical stimulation (30 Hz, pulse width 250 μ s, current 17.8 [2.9] mA) and immediately after the stimulation device was turned off. The probe was kept in the same position between the two measurements. Each Ds-t was measured by two different observers (J. Steier and J. Seymour, random order) who were blind to each other's measurements. Subjects were measured on two occasions to assess reproducibility.

Respiratory Pressures and Electromyography of the Diaphragm and Other Respiratory Muscles

These measurements were made according to the American Thoracic Society/European Respiratory Society joint statement on respiratory muscle testing²¹ and as previously described by others.²²⁻²⁷ Additional information is given in e-Appendix 1. Multiple respiratory muscle tests were performed as described elsewhere.^{21,28} Four maneuvers, maximal inspiratory effort against a closed shutter, maximum sniff, inspiration to total lung capacity, and maximum voluntary ventilation, were performed. These have been shown to produce maximal or near-maximal diaphragm activation.^{22,29} The largest rectified signals of the electromyogram (EMG) (root-mean-square of the raw data) acquired during spontaneous breathing were quantified off line (time constant 100 ms) and expressed as percentage of maximum EMG activity as derived from the maximum effort maneuvers. All recordings of EMG were sampled at 2 kHz. EMG data were amplified and filtered with a high-pass 30-Hz filter and an additional low-pass filter of 3 kHz.

Overnight Study

Eleven patients with OSA underwent the following measurements in addition to submental electrical stimulation and assessment of comfortable and maximal stimulation levels:

- Daytime fatigue symptoms were assessed by the Epworth Sleepiness Scale.³⁰
- Spirometry was performed according to international guidelines.^{31,32} Vital capacity was measured in the sitting and supine position.
- Arterialized earlobe blood was collected into a capillary tube and analyzed (Bayer Rapidlab 248; Diamond Diagnostics; Holliston, Massachusetts).
- Full polysomnography was performed using Alice 5 equipment (Respironics; Murrysville, Pennsylvania). Sleep and respiratory events were scored with standard terminology³³⁻³⁵ (additional information in e-Appendix 1).

Snoring was recorded with a microphone attached to the skin over the larynx and scored independently by two blinded investigators (J. Seymour and W. D.-C. Man.) by using a semiquantitative scale for analyzing 5-min printouts of the microphone recordings prior to, during, and after stimulation (0 = no snoring, 0.5 = very mild, 1 = mild, 2 = moderate, 3 = severe).

After positioning of the catheters and surface electrodes, the patients underwent respiratory muscle tests and went to bed;

Manuscript received October 22, 2010; revision accepted March 1, 2011.

Affiliations: From the King's College London School of Medicine, King's Health Partners (Drs Steier, Seymour, Rafferty, Jolley, and Moxham), London, England; the King's College Hospital (Mr Solomon), London, England; The State Key Laboratory of Respiratory Disease (Dr Luo), Guangzhou Medical School, Guangzhou, China; and the NIHR Respiratory Biomedical Research Unit (Drs Man and Polkey), Royal Brompton Hospital and Imperial College, London, England.

Funding/Support: This study was funded by a HEPTAGON scheme grant from the Lord Mayor, London, England (MABZAAR).

Correspondence to: Joerg Steier, MD, PhD, King's College London School of Medicine, Chest Unit, 2nd Floor Cheyne Wing, Denmark Hill, London, SE5 9RS, England; e-mail: joerg.steier@kcl.ac.uk

© 2011 American College of Chest Physicians. Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (<http://www.chestpubs.org/site/misc/reprints.xhtml>).

DOI: 10.1378/chest.10-2614

2 min of resting breathing were recorded while being awake and supine, and then lights were turned off (10:00 PM–11:00 PM). Once the patient fell asleep, the NeuroTrac Sports device was activated for 10 min during periods of upper airway occlusion, the earliest starting at 10 min after sleep onset. Current intensity was increased over a minute until the level of comfort, as recorded while awake, was reached. Stimulation was delivered during sleep stage N2 with the patient supine. The occurrence of respiratory events during stimulation, as measured by the respiratory disturbance index (RDI), was compared with the 10-min periods prior to and after stimulation. After a break of at least 1 h, stimulation was repeated to determine current intensity that would cause arousal. Stimulation intensity was slowly increased but kept below the maximum tolerated current, as documented in the initial daytime trial, until eventually an arousal from sleep was induced. This current was noted as the “arousal” intensity.

Analysis and Statistics

Results were further analyzed and plotted using SPSS, version 16 (SPSS; Chicago, Illinois) for Mac OS X (Apple; Cupertino, California) and GraphPad Prism 5.0 (GraphPad Software; San Diego, California). Following testing for normality results are given as mean (SD). A P value < 0.05 was considered significant. A repeated-measures analysis of variance was chosen to compare the results of the stimulation periods overnight to periods prior to and after stimulation, using a post hoc analysis with Bonferroni correction for multiple comparisons. Bland-Altman plots in e-Figures 1 and 2 compare repeatability and reproducibility of the ultrasonography measurements; bias and P values are stated in Table 1.³⁶

RESULTS

Ultrasonography Study

We studied 11 awake subjects with ultrasonography (seven men, aged 30 [6] years; BMI, 24.2 [3.5] kg/m²). The mean current used to stimulate genioglossus was 17.8 (2.9) mA. There was an increase in the Ds-t of 10.0% (2.8%) on the sagittal view and 9.4% (3.7%) on the coronal view, on average (all results of occasion 1 and 2 for observer 1 and 2) (Figs 2A, 2B; Table 1). The results were reproducible between observers and repeatable between occasions, without significant differences between observers or occasions (Table 1). (Additional data on the interoccasion repeatability and interobserver reproducibility are given in e-Figs 1 and 2.)

Sleep Study

We studied 11 patients with OSA (eight men, aged 51 [16] years; BMI, 42.0 [9.7] kg/m²; four ex-smokers, one active smoker, and six never smokers) during sleep (Table 2). Each patient was stimulated for at least 10 min during periods of upper airway occlusion. Two examples of the stimulation periods are given in Figures 3A and 3B. Figure 3A shows a patient in whom a relatively high current was used, whereas Figure 3B indicates the more subtle changes with

lower stimulation intensity. The mean current used during the first stimulation period was 10.1 (3.7) mA; the mean current to cause arousal from sleep during any of the succeeding stimulation periods was 14.8 (6.9) mA.

Patients' lung function and respiratory muscle strength is described in Table 2. (Additional data on individual subjects is provided in e-Tables 1–3.) Pulmonary function tests were borderline for restrictive ventilation disorder and revealed slightly low PaO₂ values. Inspiratory and expiratory muscle strength was normal or near normal.²⁷

Total sleep time was, on average, 313.4 (51.5) min, sleep efficiency was 74.6% (18.9%), and rapid eye movement-sleep time was 16.1% (6.4%) of the total sleep time. Two patients (numbers 1 and 9) had a low sleep efficiency, perhaps due to the invasiveness of the study. The nadir of oxygenation and the mean of the desaturations indicated moderately severe OSA (Table 3).




FIGURE 1. A, Frontal view of the placement of the patches. B, Lateral view of the placement of the patches (P) on the skin, half-way (arrow) between midline (M) of the chin and angle (A) of the mandible (indicated by the dotted lines) on the submental region, above a line (indicated by the solid line) cranial of the larynx (L). Patches were then connected by cables (C) to the stimulation device. Author J. Seymour pictured with permission.




FIGURE 2. Ultrasonograph of the submental region. Probe is pointing cranially as indicated by the arrows in the pictograms. A, Coronal view. B, Sagittal view. Appearance immediately before (left side of ultrasonograph) and after (right side of ultrasonograph) the electrical stimulation was turned off. Ds-t = distance skin-to-tongue surface; M = mandible; P = soft palate.

The results of the initial 10-min period of electrical transcutaneous stimulation during upper airway occlusion when asleep are shown in Table 4. (Additional data on individual subjects are provided in e-Tables 4-8). Snoring was reduced by stimulation ($P < .001$) and, although slightly higher levels of noise were observed when stimulation was turned off, it remained improved for the following 10 min. Oxygen

saturation (SaO_2) increased with stimulation ($P = .001$) and remained higher after stimulation stopped. RDI and apnea-hypopnea index (AHI) decreased significantly during stimulation ($P = .002$). In six of the 11 patients it was reduced to $< 5/\text{h}$. Inspiratory pressures were significantly diminished with electrical stimulation (esophageal pressure [Pes] and transdiaphragmatic pressure [Pdi], $P = .022$). Neural respiratory

Table 1—Results of Ultrasonography Measurements

Parameters	Occasion 1	Occasion 2	Mean Difference (Bias), mm	P Value	Observer 1	Observer 2	Mean Difference (Bias), mm	P Value
$\Delta\text{Ds-t}$ (sagittal)	5.8 (2.0) mm/ 9.8% (3.2%)	6.0 (1.1) mm/ 10.3% (2.2%)	0.15 (0.60)	.692	6.0 (1.8) mm/ 10.2% (2.9%)	5.8 (1.5) mm/ 9.9% (2.5%)	0.01 (2.6)	.733
$\Delta\text{Ds-t}$ (coronal)	5.4 (2.4) mm/ 9.6% (3.9%)	5.4 (2.0) mm/ 9.7% (3.7%)	0.19 (0.91)	.984	5.5 (2.2) mm/ 9.8% (3.8%)	5.3 (2.2) mm/ 9.5% (3.9%)	-0.20 (1.5)	.823

Values are presented as mean (SD). The results of occasions 1 and 2 are summarized in columns 2 and 3, the bias (occasion 2 - 1) in column 4, and the appropriate P value in column 5. The results of the two observers, the bias (observer 2 - 1), and P value comparing their results are given in the four right columns. An increase of around 10% in the diameter would result in an increase of approximately 21% of the cross-sectional area of the tongue, assuming an ideal fit model of a circle (circle area = πr^2). $\Delta\text{Ds-t}$ = change in the distance skin-to-tongue surface between stimulation on and off.

Table 2—Patient Characteristics, Pulmonary Function Tests, Earlobe Blood Gas Levels, Stimulation Intensity, and Respiratory Muscle Tests

Parameter	Mean (SD)
Sex, male:female	8:3
Age, y	50.7 (15.9)
BMI, kg/m ²	42.0 (9.7)
Neck circumference, cm	43.4 (3.4)
Waist/hip ratio	1.04 (0.08)
Epworth Sleepiness Scale Score, points	11.4 (6.4)
Current/stimulation, mA	10.1 (3.7)
Current/arousal, mA	14.8 (6.9)
FEV ₁ , % predicted	80.8 (15.2)
VC, % predicted	83.1 (16.3)
Change in VC, %	4.7 (3.8)
PaO ₂ , mm Hg	73.7 (7.4)
PaCO ₂ , mm Hg	40.5 (3.0)
Bicarbonate, mmol/L	25.3 (1.4)
Sniff Pdi, cm H ₂ O	122.6 (30.7)
Sniff Pes, cm H ₂ O	96.7 (23.2)
Sniff Pnasal, cm H ₂ O	82.3 (28.2)
Ptmax, cm H ₂ O	84.8 (28.8)
PEmax, cm H ₂ O	133.4 (37.0)
Cough Pgas, cm H ₂ O	230.8 (33.8)

All but the cough Pgas and PEmax test results were performed from functional residual capacity. PEmax maneuvers were performed from total lung capacity. Current stimulation: delivered over the initial 10-min period, judged as comfortable while awake. Current arousal: stimulation intensity at which arousal occurred during the second period of stimulation. Change in VC: when changing posture from sitting to supine. Cough Pgas = gastric pressure during a maximum cough; PEmax = maximum expiratory mouth pressure; Ptmax = maximum inspiratory mouth pressure; Sniff Pdi = maximum transdiaphragmatic sniff pressure; Sniff Pes = maximum esophageal sniff pressure; Sniff Pnasal = maximum nasal sniff pressure; VC = vital capacity.

drive transesophageal diaphragm electromyogram (EMGdi[%max]) was reduced during stimulation ($P < .001$) and returned, along with the RDI, to the levels prior to stimulation when the device was turned off (Fig 3, Table 4). Neural respiratory drive and Pdi during resting breathing, awake and supine, were comparable to levels during stimulation when asleep; Pes deflections remained slightly elevated (Table 4). The EEG revealed an electrical artifact during stimulation, but the first stimulation was at low levels of current and the patients did not arouse from sleep (Fig 3B).

DISCUSSION

Transcutaneous electrical stimulation of the genioglossus causes a measurable contraction of the tongue and pharyngeal muscles. This effect is reproducible and can be reliably visualized using ultrasonography. Such stimulation of the genioglossus contracts pharyngeal dilator muscles and reduces ventilatory load and neural drive in patients with OSA. Low-current transcutaneous electrical stimulation can be effec-

tively delivered intermittently overnight at levels that do not cause awakening. Overall, respiratory effort and neural respiratory drive during stimulation periods reached levels similar to those during unoccluded breathing while being awake and supine.



Clinical Significance of Findings

Given the increasing number of patients diagnosed with OSA and the observed adherence with CPAP, an alternative approach to treatment could be useful. The reduction in obstructive respiratory events in this study suggests that transcutaneous electrical stimulation of genioglossus has potential as an additional therapy in OSA. We acknowledge that those studied are a selected group of patients and a prospective randomized controlled trial is necessary to determine the usefulness of this approach.

Oliven et al¹⁷ studied electrical stimulation of the genioglossus, using fine wire electrodes that were inserted intraorally into the muscle, and measured upper airway caliber and critical occlusion pressure. Our results are consistent with their findings in that they found around one-half of the tested patients with OSA to have improved pharyngeal patency when stimulated asleep.

Miki et al¹⁹ studied six patients with OSA, placing two silver chloride ECG electrodes, 10 mm in diameter, 1 cm apart on the skin in the submental region. An apnea-triggered stimulator delivered electrical impulses (0.5 milliseconds, 50 Hz) after 5 s of apnea and ceased after airflow resumed or after 10 s. They did not confirm muscle contraction other than by observation. In their study, only two patients were tested with longer periods of stimulation. After initial tests with different stimulation frequencies, we chose a frequency of 30 Hz based on the observation that for all skeletal muscles tension is frequency dependent and stimulation frequencies of 20 to 30 Hz generate about two-thirds of maximum tension, which is well maintained for prolonged periods, whereas prolonged stimulation at 50 Hz is likely to induce fatigue.³⁷

A study by Edmonds et al²⁰ investigated transcutaneous electrical stimulation of the genioglossus in eight patients with OSA, with different findings from those of Miki et al.¹⁹ They found that transcutaneous electrical stimulation during upper airway occlusion was not effective and caused arousal. However, the intensity of stimulation in their study was relatively high. In the present study, the maximal (SD) tolerated current in normal subjects, who were highly motivated and familiar with physiologic studies, was 17.8 (2.9) mA, while being awake, compared with a range of 15 to 39.6 mA used in patients in the study of Edmonds et al.²⁰ For our overnight stimulation protocol, we used lower currents of 10.1 (3.7) mA, which did not wake the patients (Table 2). It is of interest



that when the stimulation intensity was increased the patients were aroused from sleep at 14.8 (6.9) mA, the lower end of the range of stimulation intensities used by Edmonds et al.²⁰ It may also be important that we used relatively large pads (40 mm × 40 mm) that allow stimulation of a greater muscle mass, at lower intensity, and therefore less discomfort.

Limitations to This Study

This study did not have a control group. However, we have previously published data on neural respiratory drive and inspiratory pressures in patients with obesity supine and awake.³⁸ These data are comparable to the levels of drive and inspiratory pressures that we have observed in this trial during electrical stimulation (Table 4). EMGdi supine in obese subjects was 24.7 (8.2) % max, Pes swings were 16.0 (5.0) cm H₂O, and Pdi swings were 18.2 (4.5) cm H₂O.

Transcutaneous electrical stimulation of genioglossus has clear limitations, and there are a number of unknowns when using the technique. Stimulation frequency, wave-form, ramp profile during initiation of stimulation, current intensity, and duration of stimulation need to be more fully investigated. The stimulation is nonspecific, stimulating several pharyngeal, facial, and neck muscles and nerves. Stimulation can sometimes be uncomfortable. These confounders can lead to arousals from sleep.

It could, therefore, be argued that patients woke up during stimulation and that this explains the improvement in the RDI. However, the patients did not have upper airway obstruction when supine and awake at the start of the study, and the flow signal and abdominal and chest wall movements indicated continuing obstructive apneas, of reduced extent, during stimulation. The microphone continued to record some snoring. Inspiratory pressure measurements and the diaphragm EMG demonstrated inspiratory effort that was greater than when awake. SaO₂ measurements showed ongoing, slightly reduced oxygen desaturations during stimulation (for illustration see Fig 3). Wherever EEG was of acceptable quality (Fig 3B), it confirmed that patients were asleep during and sub-

sequent to stimulation. We, therefore, concluded that the reduced level of obstruction observed during stimulation was not due to wakening. However, the electrical interference may have had an artifactual impact on EMGdi, despite using digital filters, and it is therefore necessary to carefully interpret EMGdi results.

Timing of the stimulation may have influenced the outcome, and arousal thresholds may indeed vary throughout the night. Therefore, our findings may have been influenced by the time chosen to deliver the stimulation. We have tried to deliver the stimulation in all participants as early as possible in the night, closer to sleep onset than to awakening, and arousal threshold at that time may have been higher than it would have been during early morning hours.

Two of the patients had low sleep efficiency. Although this may be caused by the invasive nature of the tests performed, it cannot be ruled out that this is due to disturbance by the stimulation, with intermittent arousal causing an underestimation of the AHI.

Frequently, patients with sleep apnea have more severe sleep-disordered breathing when supine. Therefore, nonsupine posture may alleviate upper airway obstruction and potentially have an impact on therapeutic methods. Hypothetically, stimulation in the non-supine posture may achieve the same effect at lower levels than in supine posture, but this remains to be tested.

In this study, the investigators manually turned the electrical stimulation on and off. Future studies could focus on automatic devices that could potentially be developed for domiciliary settings. Feedback mechanisms to activate the device might be best based on flow signals to treat sleep apnea and snoring, and this technique might also be used to reduce the burden of nocturnal domiciliary noise. Following the initial experience in this feasibility study we believe that a sham controlled study of patients with sleep apnea using transcutaneous electrical stimulation of the genioglossus throughout the whole of the night in comparison with a control night without stimulation, randomly assigned, is needed to assess benefit from this technique.

FIGURE 3. A, Transcutaneous stimulation of the genioglossus (stimulation sequence: off-on/on-off; 18 mA) in a patient with repetitive upper airway obstruction in sleep, shown are beginning and end of a 10-min stimulation sequence. Black bars indicate the stimulation artifact in the surface EMG signals. This patient had severe obstructive sleep apnea; there was substantial improvement during electrical stimulation. The inspiratory pressures (Pes, Pdi, both in cm H₂O) decreased, whereas there was still some upper airway obstruction during the initial period as the current was slowly increased. Abdomen and thorax indicate uncalibrated respiratory inductance plethysmography. B, Low-current stimulation in a patient with obstructive sleep apnea, beginning with 1 mA and slowly titrating up to 4 mA. A clear stimulation artifact, increasing with each upward step of current intensity, can be seen in the electromyogram of the sternocleidomastoid (EMGneck). Pes swings and neural respiratory drive decreased; there was continuous airflow with stimulation. Snoring became more continuous because of permanent airflow, and desaturations were less severe. A stimulation artifact can be observed in the EEG, which could potentially mask arousal from sleep. It is also displayed in the unfiltered raw data of the EMGdi and may have interfered with data analysis, although this signal was processed using digital filters. EMG = electromyogram; EMGchin = genioglossus surface electromyogram; EMGdi = the maximum recording of transesophageal electromyogram of the diaphragm; EMGneck = sternocleidomastoid surface electromyogram; Pdi = transdiaphragmatic pressure; Pes = esophageal pressure; Pgas = gastric pressure; SaO₂ = oxygen saturation.

Table 3—Polysomnography Data

Subject No.	TST, min	REM, min	Sleep Efficiency, %	Min SaO ₂ , %	Mean SaO ₂ Desaturation
1	146.0	25.0	61.9	87	89
2	274.0	47.5	72.9	80	85
3	380.5	73.5	88.5	67	89
4	302.0	71.5	74.8	78	82
5	366.5	107.0	87.7	84	88
6	395.5	50.5	99.9	87	90
7	389.0	61.5	88.8	72	84
8	340.0	30.5	70.2	73	83
9	126.0	10.0	28.0	88	90
10	314.5	49.0	72.1	88	91
11	413.0	40.0	75.6	76	80
Mean	313.36	51.45	74.58	80.00	86.45
SD	97.59	26.62	18.89	7.38	3.78

REM = rapid eye movement time; SaO₂ = oxygen saturation; TST = total sleep time.

Interestingly, a recent presentation of a feasibility study of hypoglossal nerve stimulation therapy to treat OSA using an implantable device (HGNS; Apnex Medical; St. Paul, Minnesota) showed comparable results to the present study with a reduction in AHI from 49.3 (18.1) to 21.6 (11.7) per h, stimulating throughout the night.³⁹ The approach of using transcutaneous stimulation could help to identify responders and nonresponders prior to implanting such devices.

Alternatives to CPAP in Sleep Apnea

In addition to CPAP, other options to treat sleep apnea include mandibular jaw advancement (dental appliances) and surgical procedures. Oral appliances need to be worn during sleep to maintain upper airway patency by increasing its diameter and reducing its collapsibility.⁴⁰⁻⁴² Their success rate is approxi-

mately 50% to 60% for partial or complete resolution of obstructive episodes, depending on the definition of complete and partial response.⁴¹⁻⁴³ Randomized controlled trials indicate that around 35% to 40% of patients with OSA have treatment failure with oral appliances.^{44,45} Best results are achieved with a custom-made mandibular advancement device.⁴⁶ The easy use of such devices with a low risk of adverse events helps to improve the compliance of patients, and they are an important treatment option in patients who are not able to cope with CPAP treatment.⁴¹⁻⁴³

In addition, there are different surgical approaches to the treatment of snoring, upper airways resistance syndrome, and OSA. The most successful surgical approach is the extirpation of hyperplastic polyps and tonsils to establish a wider lumen. Among other surgical interventions are the uvulopalatopharyngoplasty, nasal surgery, and maxillofacial surgery. However, data on outcome for these interventions are

Table 4—Parameters Awake and Asleep Prior to, During, and After Stimulation

Parameter	Awake	10 min Prestimulation	10 min During Stimulation	10 min Poststimulation
Heart rate, beats/min	84 (26)	79 (22)	69 (18)	74 (20)
Snoring, score: 0-3 points	...	2.1 (0.9)	1.4 (0.9) ^a	1.7 (0.9) ^b
SaO ₂ , %	...	91.9 (2.1)	93.2 (1.8) ^b	92.8 (2.1) ^c
RDI, events/h	...	28.1 (26.3)	10.2 (10.2) ^b	26.6 (26.0)
AHI, per h	...	26.3 (25.1)	10.0 (9.8)	25.7 (25.1)
Pes, cm H ₂ O	14.3 (6.1) ^c	23.6 (14.1)	18.3 (7.5) ^c	23.4 (11.3)
Pdi, cm H ₂ O	18.7 (6.2)	24.1 (13.5)	19.7 (7.1) ^c	24.2 (10.8)
EMGdi, μ V	25.5 (8.4)	42.9 (30.5)	26.3 (13.4) ^b	40.8 (25.7)
EMGdi, %max	15.0 (4.4)	23.8 (12.6)	15.7 (6.4) ^a	22.6 (10.4)
Current, mA	...	...	10.1 (3.7)	...

Indicated are the differences between awake and stimulation, prior to and during stimulation, and prior to and poststimulation. EMGdi and Pdi normalized to levels during wakefulness; only Pes remained elevated during stimulation. There was a significant reduction of snoring (semiquantitative scale; points) during and after stimulation. SaO₂ improved during and after stimulation. The RDI was reduced with stimulation, compared with both prior to and afterward, and inspiratory pressures (Pes, Pdi) decreased during electrical stimulation. The EMGdi indicated a decrease in neural respiratory drive during stimulation. AHI = apnea-hypopnea index; EMGdi = transesophageal diaphragm electromyogram; Pdi = transdiaphragmatic pressure; Pes = esophageal pressure; RDI = respiratory disturbance index.

^a*P* < .001.

^b*P* < .01.

^c*P* < .05.

limited in terms of quantifiable impact on physiologic parameters or long-term results, whereas such information is available for CPAP treatment. Registers and careful follow-up are recommended by critical reviewers in this field.⁴⁷

In this context, the current technique may be used in a cascade of therapeutic options to offer alternative support in the sleep laboratory to patients unable or unwilling to use CPAP. Our data indicate that some patients are more likely to benefit from this technique (six out of the 11 patients had a reduction of RDI below 5/h). We, therefore, think that this technique may be more successful in patients with mild to moderate disease and lower body mass.

CONCLUSION

In normal awake subjects, transcutaneous electrical stimulation causes contraction of the tongue muscles. In patients with OSA, snoring and sleep-disordered breathing can be improved by continuous stimulation if low current is used to avoid awakening from sleep. The feasibility and effectiveness of low-current transcutaneous genioglossus stimulation requires larger studies, alongside experimental work to determine optimal stimulation paradigms and the development of automated feedback systems.

ACKNOWLEDGMENTS

Author contributions: Dr Steier: contributed to study design, recruitment, data acquisition, data analysis, statistical analysis, and manuscript writing.

Dr Seymour: contributed to study design, recruitment, data acquisition, data analysis, and manuscript writing.

Dr Rafferty: contributed to study design, statistical analysis, and manuscript writing.

Dr Jolley: contributed to study design, recruitment, analysis, and manuscript writing.

Mr Solomon: contributed to study design, data acquisition, manuscript writing, and analysis.

Dr Luo: contributed to study design, recruitment, data acquisition, and manuscript writing.

Dr Man: contributed to study design, data analysis, and manuscript writing.

Dr Polkey: contributed to study design, recruitment, statistical analysis, manuscript writing, and revision.

Dr Moxham: contributed to study design, recruitment, statistical analysis, manuscript writing, and revision.

Financial/nonfinancial disclosures: The authors have reported to CHEST the following conflicts of interest: Dr Luo has developed the multipair catheter used for the measurement of the diaphragm EMG; no patent is pending. Drs Steier, Seymour, Rafferty, Jolley, Man, Polkey, and Moxham and Mr Solomon have reported that no potential conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Role of sponsors: The sponsor had no influence on the study design, recruitment, analysis, or manuscript writing. The funding was provided to promote independent translational research creating potentially new approaches for treatment.

Additional information: The e-Appendix, e-Figures, and e-Tables can be found in the Online Supplement at <http://chestjournal.chestpubs.org/content/140/4/998/suppl/DC1>.

REFERENCES

1. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med*. 1993;328(17):1230-1235.
2. Sullivan CE, Issa FG, Berthon-Jones M, Eves L. Reversal of obstructive sleep apnoea by continuous positive airway pressure applied through the nares. *Lancet*. 1981;1(8225):862-865.
3. Ayas NT, Mancini GBJ, Fleetham J. Does CPAP delay the development of cardiovascular disease in patients with obstructive sleep apnea hypopnea? *Thorax*. 2006;61(6):459-460.
4. Bradley TD, Logan AG, Kimoff RJ, et al; CANPAP Investigators. Continuous positive airway pressure for central sleep apnea and heart failure. *N Engl J Med*. 2005;353(19):2025-2033.
5. Senn O, Brack T, Russi EW, Bloch KE. A continuous positive airway pressure trial as a novel approach to the diagnosis of the obstructive sleep apnea syndrome. *Chest*. 2006;129(1):67-75.
6. Tawk M, Goodrich S, Kinasevitz G, Orr W. The effect of 1 week of continuous positive airway pressure treatment in obstructive sleep apnea patients with concomitant gastroesophageal reflux. *Chest*. 2006;130(4):1003-1008.
7. McArdle N, Devereux G, Heidarnejad H, Engleman HM, Mackay TW, Douglas NJ. Long-term use of CPAP therapy for sleep apnea/hypopnea syndrome. *Am J Respir Crit Care Med*. 1999;159(4 pt 1):1108-1114.
8. Remmers JE, deGroot WJ, Sauerland EK, Anch AM. Pathogenesis of upper airway occlusion during sleep. *J Appl Physiol*. 1978;44(6):931-938.
9. Bishara H, Odeh M, Schnall RP, Gavriely N, Oliven A. Electrically-activated dilator muscles reduce pharyngeal resistance in anaesthetized dogs with upper airway obstruction. *Eur Respir J*. 1995;8(9):1537-1542.
10. Oliven A, Schnall RP, Pillar G, Gavriely N, Odeh M. Sublingual electrical stimulation of the tongue during wakefulness and sleep. *Respir Physiol*. 2001;127(2-3):217-226.
11. Randerath WJ, Galetke W, Domanski U, Weitkunat R, Ruhle KH. Tongue-muscle training by intraoral electrical neurostimulation in patients with obstructive sleep apnea. *Sleep*. 2004;27(2):254-259.
12. Oliven A, Odeh M. Effect of positional changes of anatomic structures on upper airway dilating muscle shortening during electro- and chemostimulation. *J Appl Physiol*. 2006;101(3):745-751.
13. Schwartz AR, Eisele DW, Hari A, Testerman R, Erickson D, Smith PL. Electrical stimulation of the lingual musculature in obstructive sleep apnea. *J Appl Physiol*. 1996;81(2):643-652.
14. Smith PL, Eisele DW, Podszus T, et al. Electrical stimulation of upper airway musculature. *Sleep*. 1996;19(suppl 10):S284-S287.
15. Eisele DW, Smith PL, Alam DS, Schwartz AR. Direct hypoglossal nerve stimulation in obstructive sleep apnea. *Arch Otolaryngol Head Neck Surg*. 1997;123(1):57-61.
16. Oliven A, O'Hearn DJ, Boudewyns A, et al. Upper airway response to electrical stimulation of the genioglossus in obstructive sleep apnea. *J Appl Physiol*. 2003;95(5):2023-2029.
17. Oliven A, Tov N, Geitini L, et al. Effect of genioglossus contraction on pharyngeal lumen and airflow in sleep apnoea patients. *Eur Respir J*. 2007;30(4):748-758.
18. Mann EA, Burnett T, Cornell S, Ludlow CL. The effect of neuromuscular stimulation of the genioglossus on the hypopharyngeal airway. *Laryngoscope*. 2002;112(2):351-356.
19. Miki H, Hida W, Chonan T, Kikuchi Y, Takishima T. Effects of submental electrical stimulation during sleep on upper airway patency in patients with obstructive sleep apnea. *Am Rev Respir Dis*. 1989;140(5):1285-1289.

20. Edmonds LC, Daniels BK, Stanson AW, Sheedy PF III, Shepard JW Jr. The effects of transcutaneous electrical stimulation during wakefulness and sleep in patients with obstructive sleep apnea. *Am Rev Respir Dis.* 1992;146(4):1030-1036.
21. American Thoracic Society/European Respiratory Society. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med.* 2002;166(4):518-624.
22. Sinderby C, Beck J, Spahija J, Weinberg J, Grassino A. Voluntary activation of the human diaphragm in health and disease. *J Appl Physiol.* 1998;85(6):2146-2158.
23. Baydur A, Behrakis PK, Zin WA, Jaeger M, Milic-Emili J. A simple method for assessing the validity of the esophageal balloon technique. *Am Rev Respir Dis.* 1982;126(5):788-791.
24. Polkey MI, Duguet A, Luo Y, et al. Anterior magnetic phrenic nerve stimulation: laboratory and clinical evaluation. *Intensive Care Med.* 2000;26(8):1065-1075.
25. Luo YM, Harris ML, Lyall RA, Watson A, Polkey MI, Moxham J. Assessment of diaphragm paralysis with esophageal electromyography and unilateral magnetic phrenic nerve stimulation. *Eur Respir J.* 2000;15(3):596-599.
26. Luo YM, Wu HD, Tang J, et al. Neural respiratory drive during apnoeic events in obstructive sleep apnoea. *Eur Respir J.* 2008;31(3):650-657.
27. Steier J, Jolley CJ, Seymour J, et al. Sleep-disordered breathing in unilateral diaphragm paralysis or severe weakness. *Eur Respir J.* 2008;32(6):1479-1487.
28. Steier J, Kaul S, Seymour J, et al. The value of multiple tests of respiratory muscle strength. *Thorax.* 2007;62(11):975-980.
29. Luo YM, Moxham J. Measurement of neural respiratory drive in patients with COPD. *Respir Physiol Neurobiol.* 2005;146(2-3):165-174.
30. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep.* 1991;14(6):540-545.
31. Guidelines for the measurement of respiratory function. Recommendations of the British Thoracic Society and the Association of Respiratory Technicians and Physiologists. *Respir Med.* 1994;88(3):165-194.
32. Miller MR, Hankinson J, Brusasco V, et al; ATS/ERS Task Force. Standardisation of spirometry. *Eur Respir J.* 2005;26(2):319-338.
33. Iber C, Ancoli-Israel S, Chesson A, Quan SF, for the American Academy of Sleep Medicine. *The AASM Manual for Scoring of Sleep and Associated Events: Rules.* 1st ed. Westchester, Illinois: Terminology and Technical Specification; 2007.
34. Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. The Report of an American Academy of Sleep Medicine Task Force. *Sleep.* 1999;22(5):667-689.
35. Bonnet M, Carley D, Carskadon M, et al. EEG arousals: scoring rules and examples: a preliminary report from the sleep disorders atlas task force of the American sleep disorders association. *Sleep.* 1992;15(2):173-184.
36. Petrie A, Sabin C. *Medical Statistics at a Glance.* 2nd ed. Oxford, England: Blackwell Publishing Ltd; 2005.
37. Edwards RHT, Young A, Hosking GP, Jones DA. Human skeletal muscle function: description of tests and normal values. *Clin Sci Mol Med.* 1977;52(3):283-290.
38. Steier J, Jolley CJ, Seymour J, Roughton M, Polkey MI, Moxham J. Neural respiratory drive in obesity. *Thorax.* 2009;64(8):719-725.
39. Eastwood PR, Barnes M, McEvoy RD, et al. Feasibility of hypoglossal nerve stimulation therapy to treat obstructive sleep apnea [abstract]. *Am J Respir Crit Care Med.* 2010;181(1):A6767.
40. Cistulli PA, Gotsopoulos H, Marklund M, Lowe AA. Treatment of snoring and obstructive sleep apnea with mandibular repositioning appliances. *Sleep Med Rev.* 2004;8(6):443-457.
41. Ferguson KA, Cartwright R, Rogers R, Schmidt-Nowara W. Oral appliances for snoring and obstructive sleep apnea: a review. *Sleep.* 2006;29(2):244-262.
42. Chan ASL, Lee RW, Cistulli PA. Dental appliance treatment for obstructive sleep apnea. *Chest.* 2007;132(2):693-699.
43. Ayas NT, Epstein LJ. Oral appliances in the treatment of obstructive sleep apnea and snoring. *Curr Opin Pulm Med.* 1998;4(6):355-360.
44. Mehta A, Qian J, Petocz P, Darendeliler MA, Cistulli PA. A randomized, controlled study of a mandibular advancement splint for obstructive sleep apnea. *Am J Respir Crit Care Med.* 2001;163(6):1457-1461.
45. Gotsopoulos H, Chen C, Qian J, Cistulli PA. Oral appliance therapy improves symptoms in obstructive sleep apnea: a randomized, controlled trial. *Am J Respir Crit Care Med.* 2002;166(5):743-748.
46. Vanderveken OM, Devolder A, Marklund M, et al. Comparison of a custom-made and a thermoplastic oral appliance for the treatment of mild sleep apnea. *Am J Respir Crit Care Med.* 2008;178(2):197-202.
47. Pépin JL, Veale D, Mayer P, Bettiga G, Wuyam B, Lévy P. Critical analysis of the results of surgery in the treatment of snoring, upper airway resistance syndrome (UARS), and obstructive sleep apnea (OSA). *Sleep.* 1996;19(suppl 9):S90-S100.